BGC Practical Lab Notes Coupled & Ocean

Keith Lindsay, NCAR/CGD

NCAR is sponsored by the National Science Foundation





BGC in CESM

- CAM CO₂ features
 - CO₂ constituents that use LND & OCN CO₂ fluxes as surface boundary condition
 - Pass CO₂ to driver for LND & OCN flux computations
 - Couple CO₂ constituents to radiation computations

- POP Ecosystem model
 - uses MARBL library in CESM2
- CLM features covered elsewhere

Coupled BGC Compsets

- Terminology
 - BGC CO₂: what is used by surface components
 - RAD CO₂: what is used by ATM radiative code
 - Diagnostic CO₂: prescribed ATM concentrations
 - Ex: constant, read from file, 1% ramp
 - Prognostic CO₂: predicted ATM concentrations
 - atmospheric constituent computed from surface CO₂ fluxes
- B1850, BHIST
 - compset long name has BGC%BDRD
 - coupled model, BGC & RAD CO₂ are diagnostic
- use B1850, BHIST longname, replace BDRD with BPRP
 - doesn't currently have corresponding short name
 - coupled model, BGC & RAD CO₂ are prognostic
- ./query_config --compsets allactive
 - run from same directory as create_newcase

Ocean Specific BGC Compsets

- C1850ECO
 - Ocean alone, 1850 aerosols, normal year forcing
- G1850ECO
 - Ocean-Ice, 1850 aerosols, normal year forcing
- G1850ECOIAF
 - Ocean-Ice, 1850 aerosols, interannually varying forcing

./query_config --compsets pop

 run from same directory as create_newcase

Initial Conditions (IC)

- Coupled compsets
 - ICs are provided, but are not spun-up
 - Spun-up ICs will be included in CESM2.1 release
 - perhaps only for f09_g17
- Ocean Alone, Ocean-Ice
 - ICs are provided, but are not spun-up

BGC env*xml variables

- CCSM_BGC
 - Controls which CO₂ fields are exchanged between CESM components, see table at end
- CCSM_CO2_PPMV
 - Constant CO₂ ref value used in some configurations
- OCN_CO2_TYPE, LND_CO2_TYPE
 - Controls CO₂ used by ocean and land components
 - constant, prognostic, diagnostic
- OCN_TRACER_MODULES
 - Controls which ocean tracers are used
 - Ocean ecosystem model is called ecosys

Exercise

- Set up different experiments and compare resulting case directories. Do differences make sense?
- Expr 1: B1850, f19_g16
- Expr 2: Take compset long name from Expr 1 and run create_newcase with this, changing BDRD to BPRP
 - What happens when you run create_newcase?
 - follow instructions to get it working
 - don't do this for real unless you know what you're doing
- Run case.setup and preview_namelists for each case
- What changes occur when the carbon cycle is made prognostic?

POP BGC Specific Output

- ocn/hist/\$CASE.pop.h.ecosys.nday1.???-??.nc
 - Selected ocean ecosys variables at daily resolution
 - Surface flux related, productivity & functional group vertical integrals
- ocn/hist/\$CASE.pop.h.ecosys.nyear1.???.nc
 - Selected three dimensional ocean ecosys tracer budget terms

UNITS & SIGN CONVENTIONS

- CAM variables CO2, CO2_LND, CO2_OCN, CO2_FFF have units kgCO₂/kg dry air
- This is **NOT** a typical unit for carbon cycle modelers
- To convert ppm, multiply by 1e6*28.966/44
- Same quantity in different component output has
 - Different names
 - Different units
 - Different sign conventions (for fluxes)

UNITS & SIGN CONVENTIONS

Component	Variable Name	Units	Sign Convention
Atmosphere	SFCO2_LND	kgCO2/m²/s	Positive up
Land	NEE	gC/m²/s	Positive up
Atmosphere	SFCO2_OCN	kgCO2/m²/s	Positive up
Ocean	FG_CO2	mmolC/m³·cm/s nmolC/cm²/s	Positive down

CCSM_BGC settings

	CO2A	CO2B	CO2C
prog CO ₂ -> land	Y	Υ	Υ
diag CO ₂ -> land	Y	Y	Y
land CO ₂ flux -> atm		Y	Y
prog CO ₂ -> ocean	Y		Y
diag CO ₂ -> ocean	Y		Y
ocean CO ₂ flux -> atm			Y

- CO2A: land only or ocean only runs
- CO2B: atmosphere-land runs
 Ocean & Fossil Fuel CO₂ fluxes read from file
- CO2C: fully coupled runs