



Day-to-Day Variability In the Thermosphere and Ionosphere Due to Lower Atmosphere Forcing

Han-Li Liu

High Altitude Observatory

National Center for Atmospheric Research

Motivation

- Day-to-day variability in the thermosphere and ionosphere is a major concern for space weather. Understanding the causes thus key for quantifying and forecasting space weather.
- Day-to-day variability caused by lower atmosphere forcing may be comparable to those caused by geomagnetic disturbances (e.g. Rishbeth and Mendillo, 2001; Rishbeth, 2006). Though pathways not clear.
- Day-to-day variability from TIME-GCM/ECMWF study was much weaker than observed.

Methodology

- NCAR Whole Atmosphere Community Climate Model with thermosphere extension (WACCM-X): 20-year climate simulation under perpetual 2000 conditions (geomagnetically quiet, and with nudged QBO).
- SD-WACCM-X with troposphere/stratosphere constrained by MERRA reanalysis (Jan-Feb 2006).
- TIME-GCM with stratosphere/mesosphere constrained by WACCM-X/MERRA simulations results.

Numerical Model: WACCM-X

- Whole Atmosphere Community Climate Model extending from earth surface to the upper thermosphere.
- Self-consistently resolve the dynamical, physical and chemical processes (ionospheric electrodynamics under development).
- Now released as one of the atmosphere components of NCAR CESM (CESM1.0.4).
- Can be run either as a climate model or with lower/middle atmosphere nudging (specified dynamics).


Numerical Model: NCAR TIME-GCM

- NCAR TIME-GCM: Thermosphere-ionosphere-mesosphere-electrodynamics general circulation model:
 - Simulates the circulation, thermal, and compositional structures and electrodynamics from the upper stratosphere to the exosphere (30-500km) (Roble, 2000).
 - Tidal waves from the lower atmosphere specified by Global Scale Wave Model (GSWM, Hagan, 1996).
 - Lower boundary can be specified by re-analysis output (NCEP or ECMWF) or idealized planetary waves.


OR

- “Nudging” by meteorological fields (Liu et al., 2013).







Variability at Stratopause/Stratosphere: NH




Variability of Migrating Tides



Variability of Non-migrating Tides




TIME-GCM/WACCM-X/MERRA: Mean State and Tidal Variability




Liu, Yudin and Roble (2013)

Variability of Vertical Drifts


Vertical drift at 75W 11.25S



Jicamarca




Fejer and Scherliess, 2001






Variability of Zonal Drift





Zonal drift at 75W 11.25S



Zonal drift: 75W 11.25S



Variability of NmF2



Summary and Conclusion

- Large-scale waves in the upper atmosphere display large day-to-day variability from WACCM-X simulations. Compared with the stratospheric planetary waves:
 - They have a much shorter time scale.
 - They do not show clear seasonal variation.
- TIME-GCM/WACCM-X/MERRA simulation is able to produce ionospheric day-to-day variability very comparable with observations (with constant and low solar/geomagnetic activity).
- Ionospheric day-to-day variability can be caused by day-to-day variability of tides and mean wind in the thermosphere.