




NCAR CESM Working Group Meeting

Nudging Timescales and Vertical Transport in CAMChem-SD and WACCM-SD

Jessica L. Neu¹, Doug Kinnison², Jean-Francois
Lamarque², A. Sasha Glanville², Michelle L. Santee¹

¹Jet Propulsion Laboratory / California Institute of Technology, ²National Center for Atmospheric Research






Model Setup and Emissions


MLS CH₃Cl Anomalies 390K

February



Santee et al., JGR, 2013

CH₃Cl Emissions from Yoshida et al. (JGR 2006)




0.0 5.0 10.0 15.0 [Gg/grid]

Time-Varying Biomass Burning Emissions

For everything else, we use the CCMi RefC1-SD model setup and emissions. The model is nudged to MERRA meteorology

$$CH_3Cl_{bb}(m, y) = CH_3Cl_{bbYOS} * \left(\frac{CO_{bb}(m, y)}{CO_{bb}(m)} \right)$$



0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0 11.0 12.0 [Gg/grid]


The original goal of this work was to use CAMChem-SD to understand the observed variability of CH₃Cl in the upper troposphere.

We found significant differences in convective transport with 50-hour and 5-hour nudging



Observed and Modeled CH_3Cl and CO Climatologies

100 hPa CH_3Cl and CO, 50 hour nudging




October to May: maxima in UT trace gases with surface sources are seen over South America, Africa, and Indonesia

June to September: maximum abundances are found in the Asian monsoon anticyclone.

The model captures the spatial distribution of CH_3Cl and CO fairly well, but underestimates CO at 100 hPa by ~40%




Relationship Between Observed and Modeled CH_3Cl and CO over Tropical Regions



Results with 50 hour nudging

Over South America and Indonesia, the slopes of the CH_3Cl and CO correlations are significantly <1 and also much smaller than the slopes over Africa

This suggests that the model convection may be too weak over these regions *relative to the convection over Africa*






Relationship Between Observed and Modeled CH_3Cl and CO over Tropical Regions - Dependence on Nudging Timescale


50 hour nudging

CAMChem CO Clim January




Observations

MLS CO Clim January



5 hour nudging

CAMChem CO Clim January




CH_3Cl



CO

CH_3Cl



CO





Africa





S. America





Indonesia




Africa



S. America




Indonesia



5 hour nudging improves the model-measurement consistency across regions and eliminates spurious model abundances



Relationship Between Observed and Modeled CH_3Cl and CO over Tropical Regions - Dependence on Nudging Timescale





The simulation of CO anomalies is greatly improved, particularly over Africa; there is less change to CH_3Cl anomalies, but the model-measurement differences are reduced.



Convection is Strongly Dependent on the Nudging Timescale



Model Deep Convective Fluxes, 225 hPa

CAMChem DpCnvFlux Clim January



IWC Observations

MLS IWC Clim January



Convective mass fluxes over Africa and Indonesia are greatly reduced with 5 hour nudging.


MLS IWC measurements suggest comparable convective depth and intensity over South America and Africa, with stronger convection over Indonesia in January. This is more consistent with the 5 hour nudging.



Convection is Strongly Dependent on the Nudging Timescale


Model OLR

CAM-Chem OLR Clim January



OLR Observations


NOAA OLR Clim January



The model OLR is also more consistent with observations when 5-hour nudging is used.





Observed Regional Ratios of Vertical Transport Tracers





Observed Ratios of Vertical Transport Tracers over Africa and South America to Asia

OLR and 215 hPa IWC show a consistent picture that suggests that vertical transport over Asia > South America > Africa throughout the year.



The ratio of South America and Africa to Asia peaks during Mar-Apr and Nov-Dec, with minimum values during the monsoon season



At 100 hPa, IWC, CO and CH₃Cl show quite a different picture than OLR, with the strongest vertical transport over Africa during April. Vertical transport over Africa penetrates deeper into the UTLS than over the other regions




Observed vs Modeled Regional Ratios

Observations

50 Hour

5 Hour



The 5 hour nudging improves the model regional ratios compared to the observations for every tracer except CH₃Cl


In both model versions, CH₃Cl over South America has a different seasonal relationship to Asia than observed – this suggests an issue with the seasonality of South American emissions



Asian Summer Monsoon Anticyclone – 100 hPa Climatologies


July Climatologies (2005-2012) at 100 hPa

MLS v3.3



Note different scales for measurements and model

50 hour



Model trace gas concentrations in the ASM anticyclone are lower than observed (in some cases much lower) and gradients are much weaker

Model H₂O maximum values extend too far eastward and do not show a closed anticyclone signature


The model's deep convective mass flux has maximum values over the East and West coasts of India



Asian Summer Monsoon Anticyclone – 100 hPa Climatologies


July Climatologies (2005-2012) at 100 hPa

MLS v3.3




Note different scales for measurements and model

50 hour




5 hour



5 hour nudging shifts the convective region to the Bay of Bengal, but the ASM anticyclone is not as well represented, especially for CO and H₂O



Asian Summer Monsoon – July OLR






Comparison to OLR observations suggests that ASM convection is better with 5 hour nudging

So why does the 5 hour version degrade the quality of the ASM anticyclone?




WACCM Nudging Timescale: Dependence of Tropospheric Ozone Variability on the Stratosphere


Tropospheric Ozone vs Stratospheric Ozone



390 hPa Deep Convective Mass Fluxes for June CAMChem-SD 50 hour



CAMChem-SD 5 hour



SD reproduces observed stratospheric ozone variability well, but, unlike FR, does not capture the observed NH relationship between stratospheric and tropospheric ozone.

Using a 5-hour nudging timescale reduces convection over the summertime continents and greatly improves the SD performance on this metric, but degrades the overall performance in the stratosphere.



Summary and Next Steps

- ◆ The model convective mass fluxes are very sensitive to the nudging timescale. The differences in convection lead to large differences in upper tropospheric composition for some species.
- ◆ Comparison to observations tends to indicate that 5 hour nudging gives more realistic convection, which is consistent with other anecdotal findings that tropospheric composition is better simulated with 5 hour nudging.
- ◆ The ASM anticyclone, however, appears to be better simulated with a longer nudging timescale.
- ◆ Likewise, analysis of WACCM output against observations has shown that the optimal nudging timescale for the stratosphere is ~50 hours.
- ◆ These results suggest that a height-dependent nudging timescale may be necessary to accurately simulate both tropospheric and stratospheric processes (or perhaps that the suite of variables being nudged should be reconsidered).
- ◆ We are redoing analysis with the MLS averaging kernels and sampling pattern; preliminary analysis indicates the results are not strongly affected.