

Using the CLM Crop Model to assess the impacts of changes in Climate, Atmospheric, CO₂, Irrigation, Fertilizer and Geographic Distribution on Historical and Future Crop Yields

Dr. Peter Lawrence
Project Scientist


NCAR – Terrestrial Science Section
Brian O’Neill, Sam Levis, Xiaolin Ren and many others



Gridcell



CLM 5 Crop subgrid tiling



Landunit



Vegetated



Lake



Urban




Glacier



Crop

Crop Model




Land Use Change



Planting



Leaf emergence



Irrig / Fertilize



Grain fill



Harvest



Unirrig



Irrig



Unirrig



Irrig



Crop1



Crop1



Crop2



Crop2 ...

CLM Crop – Crop Simulations

We would like to evaluate agriculture in CLM Crop with CESM forcing from the Historical, RCP 4.5 and RCP 8.5 time periods

Problems with CLM4.5 Crop:

- Fixed Crop distribution as crop land unit can not be changed through time like PFTs
- Only simulates Cotton, Corn, Rice, Soy, Sugarcane, Wheat (temperate/tropical)
- Nitrogen fertilizer is fixed by crop type independent of region or time period based on North American application rates
- Has fixed irrigation area based on crop distribution

CLM Crop – Idealized Crop Simulations

Idealized Crop simulations are suite of CESM simulation with transient CESM forcing from the Historical, RCP 4.5 and RCP 8.5 time periods that can be combined with cropping and management scenario

Globally simulate in CLM Crop for every land grid cell:

- Cotton, Corn, Rice, Soy, Sugarcane, Wheat
(temperate/tropical)

With Management:

- N Fertilizer and Irrigated
- N Fertilizer and Rainfed
- No N Fertilizer and Irrigated
- No N Fertilizer and Rainfed



With Atmospheric Forcing as:

- Transient Climate and CO₂
- Transient Climate and Constant CO₂
- Constant Climate and Constant CO₂



CLM Crop – Scenario Simulations

Crop Area time series CMIP5 / other scenario

CESM Climate / CO₂ Scenario – Hist/RCP4.5/RCP8.5



Crop Type / Irrigation time series
Portmann et al. (2010)



Global CLM Crop Simulation

- Scenario
- Crop
- Year
- Area
- Yield
- Fertilizer
- Irrigation
- Temp.
- Precip.
- Solar
- CO₂


Global Idealized CLM Crop Simulation Database

For Historical, RCP4.5 and RCP8.5 time series and for each crop


Rainfed / Irrigated	N Fertilizer	CO2 Concentration	Climate
Irrigated	UNFAO Fertilizer	Transient	Transient
Rainfed	UNFAO Fertilizer	Transient	Transient
Irrigated	No/Const Fertilizer	Transient	Transient
Rainfed	No/Const Fertilizer	Transient	Transient
Irrigated	UNFAO Fertilizer	Constant	Transient
Rainfed	UNFAO Fertilizer	Constant	Transient
Irrigated	No/Const Fertilizer	Constant	Transient
Rainfed	No/Const Fertilizer	Constant	Transient
Irrigated	UNFAO Fertilizer	Constant	Constant
Rainfed	UNFAO Fertilizer	Constant	Constant
Irrigated	No/Const Fertilizer	Constant	Constant
Rainfed	No/Const Fertilizer	Constant	Constant

Global Historical CMIP5 (All Crops) Area, Yields, N Fert


Historical Global Crop Area (millions hectares)




Historical Global Production (millions tonnes)



Historical Global Crop Yield (tonnes/ha)




Historical Global N Fertilizer (kilograms/ha)




Area Effect	Climate / CO2	N Fert Effect	Irrig Effect	Total Effect
+77.2%	+5.8%	+65.6%	+9.5%	+226.8%

Global Historical CMIP5 (All Crop) Crop Production


Historical Crop area effect (2000 - 1905) Yield 10^3 Tonnes




Historical Climate and CO2 effect (2000) Yield 10^3 Tonnes



Historical N Fertilizer effect (2000) Yield 10^3 Tonnes




Historical Combined effect (2000 - 1905) Yield 10^3 Tonnes




Global Historical CMIP5 – UNFAO Area, Yields, N Fert


Historical Global Crop Area (millions hectares)




Historical Global Production (millions tonnes)



Historical Global Crop Yield (tonnes/ha)





Historical Global N Fertilizer (kilograms/ha)



Area Effect	Climate / CO2	N Fert Effect	Irrig Effect	Total Effect
+77.2%	+5.8%	+65.6%	+9.5%	+226.8%

*UNFAO only using Yields for CLM Crop Types over All Crop Areas


Global Historical CMIP5 UNFAO Yields by Crop



*UNFAO Sugar Cane scaled by 0.12 for sugar from cane


CLM Crop – UNFAO Regional N Fertilizer Application

Regional Current Day CLM Crop N Fertilizer compare to UN FAO estimates



CLM Crop – N Fertilizer Application

Current Day and Projected UN FAO N Fertilizer Application Rates (1995 – 2050)




Historical CLM Crop and UN/FAO - Yield Summary


1. CLM Crop represents the world with six crops with climate varieties as well as rainfed and irrigated management.
2. Historical Crop Area increased from 1901 – 2005 by 685 million hectares (83%) and Industrial Fertilizer increased from 0 to 77 kg/ha
3. Crop Area increases resulted in an increase in Production of 77% with Fertilizer another 66% and irrigation 9.5%. CO₂ only added another 6%
4. Compared to UNFAO, CLM Crop has a similar Production and Yield amounts for the crops simulated for current day
5. UNFAO increases in Production and Yield occur slightly more quickly than CLM Crop (1961 – 2005) possibly due to differences in crop varieties and management such as planting density and irrigation.
6. Default N Fertilizer for CLM Crop are not applicable for much of the world and need to be changed to transient regional values

Global CMIP5 RCP 8.5 (All Crops) Area, Yield, N


RCP 8.5 Global Crop Area (millions hectares)




RCP 8.5 Global Crop Production (millions tonnes)



RCP 8.5 Global Crop Yield (tonnes/ha)




RCP 8.5 Global N Fertilizer (kilograms/ha)




Area Effect	Clim Effect	CO2 Effect	N Fert Effect	Irrig Effect	Total Effect
+21.2%	-23.4%	+43.3%	+39.6%	+8.0%	+64.6%

Global CMIP5 RCP 8.5 Analysis (All Crop) Production


RCP8.5 Crop Area effect (2095 - 2010) Yield 10^3 Tonnes





RCP8.5 Climate (Const CO2) effect (2095) Yield 10^3 Tonnes




RCP8.5 CO2 Fertilization effect (2095) Yield 10^3 Tonnes



RCP8.5 N Fertilizer effect (2095) Yield 10^3 Tonnes



Global CMIP5 RCP 8.5 Analysis (All Crop) Production



Future CLM Crop – Yield Summary

1. RCP 8.5 has an increases in: cropping area of 280 million hectares (18%); average land temperature of +4°C; CO₂ to 930 ppm (150%); and Fertilizer to 117 kg/ha (60%)
2. RCP 8.5 CLM Crop results in changes in Global Production of:

Area Effect	Clim Effect	CO2 Effect	N Fert Effect	Irrig Effect	Total Effect
+21.2%	-23.4%	+43.3%	+39.6%	+8.0%	64.6%

3. Future climate impacts may be under estimated as CLM Crop does not properly account for damage from Heat Waves, Droughts, Ozone, Insects, Diseases, and Floods.
4. Currently CLM Crop responds to temperature, vapor pressure, soil moisture and light to limit photosynthesis which then impacts grain production, but this is only one element impacting yield.

CLM Crop Simulations – Conclusions

1. CLM Crop allows us to investigate changes in Agricultural Production as climate and CO₂ change in combination with cropping area and management
2. Historical increases in cropping area result in a +77% gain with fertilizer and irrigation adding an additional +75% gain. CO₂ results in only another +6% gain.
3. RCP 8.5 crop expansion results in a +21% gain which is offset by climate losses of -23%. CO₂ results in a +43% gain which combines with a fertilizer and irrigation gain of +48% to leave a net +65% gain in Production for the scenario
4. Historically differences from UNFAO may be due no management to change varieties and planting densities. Future impacts do not properly account for damage from Heat Waves, Droughts, Ozone Insects, Disease, and Floods. All of these are current research

Thanks – Questions?